

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 983 990 A2**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
08.03.2000 Bulletin 2000/10

(21) Application number: 99810685.0

(22) Date of filing: 29.07.1999

(51) Int Cl.7: **C07C 49/84, C07C 69/145,
C07C 69/712, C07C 69/74,
C07C 69/675, C07C 69/24,
C07C 69/96, C07C 229/22,
C07C 329/06, C11D 3/50,
C11D 3/00, A61K 7/46**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 04.09.1998 EP 98810883

(71) Applicant: **Glaxo S.A. (International) S.A.**
1214 Vernier-Genève (CH)

(72) Inventors:
• **Anderson, Denise Dr.**
8032 Zürich (CH)
• **Frater, Georg Dr.**
8400 Winterthur (CH)

(74) Representative: **Patentanwälte
Schaad, Balass, Menzi & Partner AG
Dufourstrasse 101
Postfach
8034 Zürich (CH)**

(54) **Ketones useful as precursors for organoleptic compounds**

(57) The ketones of the formula I are useful for the delivery of organoleptic compounds, especially for fla-

vours, fragrances and masking agents and antimicrobial compounds.

EP 0 983 990 A2

Description

[0001] The present invention relates to new ketones. These ketones are useful as precursors for organoleptic compounds, especially for fragrances and masking agents and antimicrobial compounds.

[0002] A principal strategy currently employed in imparting odours to consumer products is the admixing of the fragrance directly into the product. There are, however, several drawbacks to this strategy. The fragrance material can be too volatile and/or too soluble, resulting in fragrance loss during manufacturing, storage, and use. Many fragrance materials are also unstable over time. This again results in loss during storage.

[0003] In many consumer products it is desirable for the fragrance to be released slowly over time. Microencapsulation and inclusion complexes with cyclodextrins have been used to help decrease volatility, improve stability and provide slow-release properties. However, these methods are for a number of reasons often not successful. In addition, cyclodextrins can be too expensive.


[0004] Fragrance precursors for scenting fabrics being washed in the presence of a lipase-containing detergent are described in WO 95/04809. The fragrance precursors contained in the detergent and/or in the softener are cleaved by the lipase and a single odoriferous compound, either an odoriferous alcohol or ketone is yielded. Thereby a prolonged scenting effect on the fabric is obtained.

[0005] EP-A-0 816 322 describes fragrance precursor compositions containing carbonic acid esters or thiocarbonic acid esters.

[0006] These compositions are used in cosmetic products or in laundry products. The precursors produce fragrances upon contacting the skin or when used in the presence of lipases, e.g. as used in detergents, thus providing a prolongation of the fabric scenting effect.

[0007] One object of the present invention is to provide new precursors for compounds with different activities. A further object of the invention is to provide new compounds which are stable in a variety of applications including detergents, fabric softeners, cleaning products, personal care products, cosmetics, and sun protection products. A further object of the invention is to provide new compounds which can be cleaved by light, heat, hydrolysis and/or enzymes into one or more organoleptic compounds.

[0008] The present invention relates to new ketones of the formula I



wherein

Y is an optionally substituted alkyl, cycloalkyl, cycloalkylalkyl, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, alkylloxycarbonyl, cycloalkylloxycarbonyl, aryloxycarbonyl, amide, oximinocarbonyl, alkylthiocarbonyl, cycloalkylthiocarbonyl, whereby all alkyl groups may be straight or branched and all alkyl and cycloalkyl groups may be saturated or unsaturated,

R¹ is hydrogen or a C₁₋₆ alkyl group which may be substituted, saturated or unsaturated, straight or branched,

A is a chromophoric aromatic or heteroaromatic ring or ring system,

A may carry 1 to 8 substituents selected from hydrogen, hydroxy, mercapto, amino, acyl, ester, esterified hydroxy group, cyano and nitro; alkyl, alkoxy, alkylthio and arylalkoxy which may be substituted, saturated or unsaturated, straight or branched; aryl and heteroaryl which may be substituted; one of the substituents may be a polymeric group, and

n is an integer

with the exception of
2-ethoxy-1-phenyl-ethanone,

hydroxy-acetic acid 2-oxo-2-phenyl-ethyl ester,
acetic acid 1-methyl-2-oxo-2-phenyl-ethyl ester, and
p-methoxyphenacyl acetate.


[0009] All groups containing the term alkyl include also the unsaturated residues, e.g. alkenyl, alkynyl, cycloalkenyl etc.

[0010] The compounds of the formula I are not limited to any particular stereoisomers, all possible stereoisomers (E/Z isomers, enantiomers, diastereomers) and all mixtures are thus included within the scope of the invention.

[0011] The compounds of the formula I are mostly or nearly odourless at room temperature, atmospheric conditions and about 20 to 100 % relative humidity. However, under activating conditions, they are cleaved and one or more active compounds with organoleptic properties are generated.

[0012] The activating conditions which lead to cleavage and the desired active compounds comprise the presence of light, such as room light and sunlight, sunlight being preferred. Additional activating conditions may be heat, hydrolysis and/or enzymes.

[0013] The compounds of the formula I yield after cleavage an aryl or heteroaryl ketone of the formula II



[0014] Preferred compounds are those which yield an organoleptic aryl ketone.

[0015] The compounds of the formula I, wherein Y is an alkyl, cycloalkyl or cycloalkylalkyl group, are cleaved into an aryl or heteroaryl ketone and an aldehyde or ketone, at least one thereof is organoleptic. Examples of organoleptic aryl ketones as well as of ketones and aldehydes including the preferred and more preferred ones are listed below. Examples of organoleptic heteroaryl ketones are 2-acetylpyrazine, 2-acetyl-3-methylpyrazine, 2-acetyl-5-ethylpyrazine, 2-acetylpyridine and the acetylpyridines which are disclosed in US patent 5,214,027.

[0016] The compounds of the formula I, wherein Y is an alkyl-, cycloalkyl- or aryloxy-carbonyl group, are cleaved into an aryl or heteroaryl ketone and an alcohol, phenol, aldehyde or ketone, at least one thereof is organoleptic. Examples of alcohols and phenols including the preferred and more preferred ones are listed below.

[0017] The term phenol means an aromatic hydroxy compound wherein the hydroxy group is attached directly to the benzene ring.

[0018] The compounds of the formula I, wherein Y is an alkyl-, cycloalkyl- or arylcarbonyl group, yield after cleavage an aryl or heteroaryl ketone and an acid or a lactone, the aryl or heteroaryl ketone and/or the lactone being organoleptic. When the alkyl residue of the alkylcarbonyl group is substituted in alpha- or beta-position with a hydroxy group the released acid may be active as a peeling ingredient in cosmetics. Lactones may be formed when the alkyl residue is substituted in gamma- or delta-position with a hydroxy group or with the group O-CO-OR² or O-CO-R², wherein R² is an alkyl, aryl or aralkyl group which may be branched, unsaturated and/or substituted. Preferred compounds are those wherein R² is a C₁₋₃₀ alkyl group or a phenyl C₁₋₁₀ alkyl group. Compounds of the formula I with the group O-CO-OR² yield after cleavage besides the aryl or heteroaryl ketone and lactone an alcohol, phenol, aldehyde or ketone. The three obtained compounds can be organoleptic.

[0019] In the compounds of the formula I, wherein Y is an alkylcarbonyl group, the alkyl group may be substituted e.g. with an oxo group. These compounds yield after cleavage an aryl or heteroaryl ketone and a ketone, at least one thereof is organoleptic. Preferred compounds are those wherein the alkyl group is substituted in beta-position to the carbonyl group with the oxo group or in delta-position, in the latter case the oxo group may be situated in an optionally unsaturated cycloalkyl radical which may be substituted.

[0020] The compounds of the formula I, wherein Y is an amide, i.e. a N-substituted aminocarbonyl group, are cleaved into an aryl or heteroaryl ketone and an amine, at least one thereof is organoleptic. Examples of organoleptic amines including the preferred are listed below.

[0021] The compounds of the formula I, wherein Y is an alkylthiocarbonyl or cycloalkylthiocarbonyl group, are cleaved into an aryl or heteroaryl ketone and a thiol, at least one thereof is organoleptic. Examples of organoleptic thiols are 3,7-dimethyl-1-thio-octa-2,6-diene, 2-(1-mercapto-1-methylethyl)-5-methylcyclohexanone and 2-methoxy-2-methylbutan-2-thiol.

[0022] The compounds of the formula I, wherein Y is an oximinocarbonyl group, i.e. -CO-O-N=CR³R⁴, wherein R³ and R⁴ may be e.g. an optionally substituted, straight or branched, saturated or unsaturated alkyl group, are cleaved into an aryl or heteroaryl ketone and an oxime, at least one thereof is organoleptic.

[0023] The present invention includes also compounds of the formula I which are cleaved into an optionally organoleptic

noleptic aryl or heteroaryl ketone and an organoleptic nitrile. In this case Y stands for an alkyl- or alkenylcarbonyl group which is substituted with a cyano group in alpha- and/or gamma-position to the carbonyl group, with the proviso that the chain between the carbonyl group and the cyano group has a double bond in beta-gamma-position to the carbonyl group when the cyano group stands in gamma-position. The alkyl or alkenyl group may be branched and/or substituted with further residues which may together form a ring system. Examples of organoleptic nitriles including the preferred ones are listed below.

[0024] In the compounds of the formula I A is a chromophoric aromatic or heteroaromatic ring or ring system. The term ring system includes mono- and polycyclic rings, which may be connected either directly or by a bridge, e.g. an optionally unsaturated alkylene group, oxygen etc., as well as fused rings. The heteroaromatic rings include five- and six-membered rings.

[0025] In most of the cases the obtained aryl or heteroaryl ketone is organoleptic. When one of the substituents of A represents a polymeric group or when A represents a fused aromatic or heteroaromatic ring system with more than two rings, the compounds of formula I release on cleavage a non-organoleptic aryl or heteroaryl ketone. In these cases O-Y in formula I represents a group which after cleavage yields at least an organoleptic aldehyde, an organoleptic ketone, an organoleptic alcohol or phenol, an organoleptic lactone, an organoleptic amine, an organoleptic thiol, an organoleptic oxime or an organoleptic nitrile, as described above.

[0026] Suitable polymeric groups are divinylbenzene-based polymers, ethyleneglycol dimethacrylate-based polymers, acrylic acid/maleic acid copolymers, acrylic acid homopolymers, polysiloxanes, alginic acid, carboxymethyl cellulose, xanthan gum and so on.

[0027] Since the compounds of the formula I, after cleavage, provide at least one substance of the group aryl or heteroaryl ketones, aldehydes, ketones, alcohols, phenols, amines, thiols, oximes, nitriles and lactones having organoleptic activity they permit the development of useful consumer products with enhanced organoleptic properties. The obtained organoleptic substances are useful as fragrances and masking agents and antimicrobial agents. Therefore, the present invention also relates to the use of all compounds of formula I as precursors for organoleptic compounds, in particular fragrances.

[0028] The compounds of the formula I can act as fragrance precursors in laundry products, such as detergents and fabric softeners, cleaning compositions, such as hard surface and all-purpose cleaners, personal care products, cosmetics or sun protection products. They can also act as precursors for odour masking agents in the same products as the fragrance precursors. They also can act as precursors for antimicrobial agents. The fragrance precursors and the precursors for odour masking agents may be used individually in an amount effective to enhance or to mask the characteristic odour of a material. More commonly the compounds are mixed with other fragrance components in an amount sufficient to provide the desired odour characteristics.

[0029] Due to the in situ generation of the active compounds the desired effect is prolonged and the substantivity on different substrates is enhanced. If two or more active compounds are provided, they can be generated, depending on the precursor simultaneously or successively. Further, the precursors of the invention provide slow release of the active compounds.

[0030] One preferred group of the ketones of the formula I are those wherein Y is an alkyl group. The alkyl group may contain a C₁₋₃₀ alkyl moiety, which may be unsaturated, branched and/or substituted by different groups as shown in the listed aldehydes and ketones below.

[0031] Preferred compounds of the formula I are those, wherein A is one aromatic ring and two of its substituents form a second ring, preferably a six-membered aromatic ring or a five- or six-membered aliphatic ring. Within this group of compounds it is most preferred that R¹ is hydrogen and the other substituents of the aromatic ring are hydrogen or one or more methyl groups.

[0032] Preferred compounds of the formula I are also those, wherein R¹ is hydrogen or methyl. Preferably A carries no more than three substituents, especially methyl groups. A further preferred group of compounds of the formula I are those, wherein R¹ is hydrogen and A is one aromatic ring, which is unsubstituted or substituted with one methoxy group, preferably in the para-position.

[0033] When Y is an alkyloxycarbonyl group, C₁₋₃₀ alkyloxycarbonyl groups are preferred which may be unsaturated, branched and/or substituted by different groups as shown in the listed alcohols, aldehydes and ketones below. The aryloxycarbonyl group is preferably a phenyloxycarbonyl group which may be substituted as shown in the listed phenols. The cycloalkyloxycarbonyl group is preferably a C₃₋₈ cycloalkyloxycarbonyl group which may be unsaturated and/or substituted.

[0034] When Y is an alkylcarbonyl group, C₁₋₃₀ alkyl groups are preferred. They may be branched, unsaturated and/or substituted, e.g. by naphthol, phenoxy, naphthyloxy, hydroxy, amino, pyrocatechol, the group O-CO-OR² or O-CO-R², as stated above. When Y is a cycloalkylcarbonyl group, C₃₋₈ cycloalkyl moieties are preferred which can be unsaturated and/or substituted. When Y is an arylcarbonyl group, the phenylcarbonyl groups are preferred which may be substituted.

[0035] When Y is an amide (N-substituted aminocarbonyl) group a wide variety of non-organoleptic amines can be


obtained by cleavage. For example, a list of suitable primary and secondary cosmetic amines can be found in the Cosmetic Ingredient Handbook edited by Joanne M. Nikitakis. Suitable surfactant amines can be found, for example, in Surfactants Europe edited by Gordon L. Hollis.

[0036] Preferred N-substituted aminocarbonyl groups are those which yield after cleavage the organoleptic amines listed below.

[0037] It is preferred that n is 1 or 2.

[0038] All groups for which Y stands may be substituted with one or more substituents selected from aryl; cycloalkyl; cycloalkenyl; oxo; oxy; cyano; hydroxy; hydroxy being esterified with a carboxylic acid, a carbonic acid ester, a N,N-substituted carbamic acid or an oximino carbonic acid; amino substituted with oximinocarbonyl; an esterified carboxylic group; and a carboxylic group esterified with an oxime.

[0039] The groups for which Y stands may also be substituted with the following substituents: A-CO-CHR¹-O-, A-CO-CHR¹-O-CO-, A-CO-CHR¹-O-alkoxy or A-CO-CHR¹-O-CO-alkoxy. The following formula III shows one example of this kind of compounds, wherein Y is an alkyl group substituted with A-CO-CHR¹-O-:



wherein m is an integer.

[0040] Compounds of the formula I may generate the following organoleptic aryl ketones:

- 4-methoxyphenyl-ethanone*
- 1-[6-(1,1-dimethylethyl)-2,3-dihydro-1,1-dimethyl-1H-inden-4-yl]-ethanone
- 1-(5,6,7,8-tetrahydro-3,5,5,6,8,8-hexamethyl-2-naphthalenyl)-ethanone*
- 2-(1-methylethyl)-indanone*
- 4-tert-butyl-3,5-dinitro-2,6-dimethyl-acetophenone
- 1,6,7,8-tetrahydro-1,4,6,6,8,8-hexamethyl-as-indacen-3(2H)-one*
- 1-(2-naphthalenyl)-ethanone*
- 1-(2,3-dihydro-1,1,2,3,3,6-hexamethyl-1H-inden-5-yl)-ethanone
- 1-[2,3-dihydro-1,1,2,6-tetramethyl-3-(1-methylethyl)-1H-inden-5-yl]-ethanone
- 3-methyl-1-(4-methylphenyl)-4-hexen-1-one
- 5-acetyl-1,1,2,3,3-pentamethylindane
- 1-phenylpropanone
- acetophenone*
- 2,4-dimethylphenyl-ethanone*
- 1-[4-(1,1-dimethylethyl)-2,6-dimethylphenyl]-ethanone
- 1-(hexahydrodimethyl-1H-benzindeny)-ethanone*
- 1-(5,6,7,8-tetrahydro-2-naphthalenyl)-ethanone
- 1-phenyl-4-penten-1-one
- 1-(5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl)-ethanone
- 1-(3-ethyl-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-ethanone

whereby * indicates the preferred aryl ketones.

[0041] Compounds of the formula I may generate the following ketones:

- 2,5-dimethyl-oct-2-en-6-one**
- 4-(2,6,6-trimethylcyclohex-1-en-1-yl)butan-2-one**
- 4-(2,6,6-trimethylcyclohex-2-en-1-yl)butan-2-one**
- 2-methyl-5-(1-methylethenyl)-cyclohex-2-en-1-one*
- 1-(4-hydroxyphenyl)-butan-3-one**
- 4-benzo-1,3-dioxo-5-yl-but-2-one**
- 2-heptyl-cyclopentanone
- nonan-2-one*
- octan-2-one*

2,2,6,10-tetramethyltricyclo-[5.4.0.0(6,10)]-undecan-4-one
 heptan-2-one*
 undecan-2-one*
 benzylacetone*
 5 butan-2-one*
 1,2,3,5,6,7-hexahydro-1,1,2,3,3-pentamethyl-4H-inden-4-one*
 6-methyl-hept-5-en-2-one*
 2-(butan-2-yl)-cyclohexanone*
 2-hexyl-cyclopent-2-en-1-one*
 10 2-(1-methylethyl)-5-methyl-cyclohexanone*
 2-(2-methylethyl)-5-methyl-cyclohexanone*
 3-methyl-cyclopentadecanone
 4-(1,1-dimethylpropyl)-cyclohexanone*
 6,10-dimethyl-undeca-5,9-dien-2-one*
 15 3-oxo-2-pentyl-cyclopentane-acetic acid methyl ester**
 1-(1,2,3,4,5,6,7,8-octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)-ethanone*
 3-methyl-5-propyl-cyclohex-2-en-1-one*
 1-(2-cyclohexen)-2,4,4-trimethyl-but-2-enone*
 carvone**
 20 2-hexyl-cyclo-pent-2-en-1-one**
 2-pentyl-cyclopent-2-en-1-one
 3-methyl-2-pentyl-cyclopent-2-en-1-one**
 2-hexylidene-cyclopentanone*
 3,5-diethyl-5,6-dimethyl-2-cyclohexenone*
 25 4,4a,5,6,7,8-hexahydro-6-isopropenyl-4,4a-dimethyl-2(3H)-naphthalenone**
 3-methyl-6-propylidenecyclohexanone*
 4-(1-methylethyl)-cyclohex-2-en-1-one
 (E)-oct-3-en-2-one
 1-(2,3,4,7,8,8A-hexahydro-3,6,8,8-tetramethyl-1H-3A,7-methanoazulen-5-yl)-ethanone*
 30 2-hydroxy-3,5-dimethyl-cyclopent-2-ene-1-one*
 1-(3,3-dimethyl-1-cyclohexen-1-yl)ethanone*
 1-(2,4,6-trimethylcyclohex-3-en-1-yl)-but-1-en-3-one acetylisolongifolene
 2-(3-methylbut-2-en-1-yl)-3-methyl-cyclopent-2-en-1-one
 3-methyl-5-(2,2,3-trimethylcyclopent-3-en-1-yl)pent-3-en-2-one*
 35 5-butylidene-2,2,4-trimethylcyclopentanone
 1,2,3,5,6,7-hexahydro-1,1,2,3,3-pentamethyl-4H-inden-4-one**
 3-methyl-5-propyl-cyclohex-2-en-1-one**
 4,4a,5,6,7,8-hexahydro-6-isopropyl-2(3H)-naphthalenone
 3,5,5-trimethyl-cyclohex-2-en-1,4-dione*
 40 (E)-5-methyl-2-hepten-4-one
 acetyl diisoamylene**
 dec-3-en-2-one
 2-ethyl-3,6,6-trimethylcyclohex-2-enyl-but-2-en-1-one
 1-(5,5-dimethyl-1(6)-cyclohexen-1-yl)-4-penten-1-one**
 45 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-but-2-en-1-one**
 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-but-2-en-1-one**
 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-but-2-en-1-one**
 2,4,4,5,5-pentamethyl-1-cyclopentene-1-yl-ethanone*

50 whereby * indicates the preferred ketones and ** indicate the more preferred ketones.

[0042] Compounds of the formula I may generate the following aldehydes:

2,6,10-trimethylundec-9-enal*
 1,2,3,4,5,6,7,8-octahydro-8,8-dimethyl-2-naphthalenecarboxaldehyde
 55 tridecanal
 2-[4-(1-methylethyl)phenyl]-ethanal
 2,4-dimethyl-cyclohex-3-ene-1-carboxaldehyde*
 4-carboxaldehyde-1,3,5-trimethyl-cyclohex-1-ene*

	1-carboxaldehyde-2,4-dimethyl-cyclohex-3-ene*
	1-carboxaldehyde-4-(4-hydroxy-4-methylpentyl)-cyclohex-3-ene*
	hex-2-enal*
	3,5,5-trimethyl-hexanal
5	heptanal*
	2,6-dimethyl-hept-5-eneal*
	decanal**
	dec-9-enal
	dec-4-en-1-al
10	2-methyldecanal*
	undec-10-ene-1-al**
	undecanal*
	dodecanal**
	2-methyl-undecanal**
15	tridecanal
	2-tridecanal
	octanal**
	nonanal*
	3,5,5-trimethylhexanal
20	2-nonenal
	undec-9-enal**
	2-phenyl-propanal*
	4-methyl-phenyl acetaldehyde*
	3,7-dimethyl-octanal*
25	dihydrofarnesal**
	7-hydroxy-3,7-dimethyl-octanal*
	2,6-dimethyl-oct-5-ene-1-al
	2-(4-(1-methylethyl)phenyl)-ethanal*
	3-(3-isopropyl-phenyl)-butanal**
30	2-(3,7-dimethyl-oct-6-en-oxy)-ethanal
	1-carboxaldehyde-4-(4-methyl-3-penten-1-yl)-cyclohex-3-ene*
	2,3,5,5-tetramethyl-hexanal
	longifolic aldehyde
	2-methyl-4-(2,6,6-trimethylcyclohex-2-en-1-yl)-butanal*
35	2-methyl-3-(4-tert-butylphenyl)propanal**
	4-(1,1-dimethyl-ethyl)-benzenepropanal*
	2-[4-(1-methyl-ethyl)phenyl]-propanal
	alpha-methyl-1,3-benzodioxole-5-propanal*
	3,7-dimethyl-oct-6-en-1-al*
40	2-methyl-3-(p-isopropylphenyl)-propionaldehyde*
	4-(4-hydroxy-4-methyl-pentyl)-cyclohex-3-en-1-carboxaldehyde**
	alpha-methyl-1,3-benzodioxole-5-propanal*
	1-carboxaldehyde-4-(1,1-dimethylethyl)-cyclohexane
	4-(octahydro-4,7-methano-5H-inden-5-ylidene)-butanal
45	[(3,7-dimethyl-6-octenyl)oxy]-acetaldehyde**
	3,7-dimethyl-oct-2,6-dien-1-al*
	nonadienal*
	2,4-dimethyl-2,6-heptadienal
	trans-dec-2-en-1-al*
50	2,4-diethyl-hep-2,6-dien-1-al*
	dodec-2-en-1-al*
	3,7-dimethyl-oct-2,6-dien-1-al*
	2,4-diethyl-hepta-2,6-dienal
	3,7-dimethyl-nona-2,6-dien-1-al*
55	3-propyl-2-hepten-1-al
	1-carboxaldehyde-4-(prop-2-en-2-yl)-cyclohex-1-ene

whereby * indicates the preferred aldehydes and ** indicate the more preferred aldehydes.

[0043] Compounds of the formula I may generate the following alcohols and phenols:

amyl alcohol
 hexyl alcohol*
 5 2-hexyl alcohol*
 heptyl alcohol*
 octyl alcohol*
 nonyl alcohol*
 decyl alcohol*
 10 undecyl alcohol*
 lauryl alcohol*
 myristic alcohol
 3-methyl-but-2-en-1-ol*
 3-methyl-1-pentanol
 15 cis-3-hexenol*
 cis-4-hexenol*
 3,5,5-trimethyl-hexanol
 3,4,5,6,6-pentamethylheptan-2-ol*
 citronellol*
 20 geraniol*
 oct-1-en-3-ol
 2,5,7-trimethyl-octan-3-ol
 2-cis-3,7-dimethyl-2,6-octadien-1-ol
 6-ethyl-3-methyl-5-octen-1-ol*
 25 3,7-dimethyl-oct-3,6-dienol*
 3,7-dimethyloctanol*
 7-methoxy-3,7-dimethyl-octan-2-ol*
 cis-6-nonenol*
 5-ethyl-2-nonanol
 30 6,8-dimethyl-2-nonanol*
 2,2,8-trimethyl-7(8)-nonene-3-ol
 nona-2,6-dien-1-ol
 4-methyl-3-decen-5-ol*
 dec-9-en-1-ol
 35 benzylalcohol
 2-methyl-undecanol
 10-undecen-1-ol
 1-phenyl-ethanol*
 2-phenyl-ethanol*
 40 2-methyl-3-phenyl-3-propenol
 2-phenyl-propanol*
 3-phenyl-propanol*
 4-phenyl-2-butanol
 2-methyl-5-phenyl pentanol*
 45 2-methyl-4-phenyl-pentanol*
 3-methyl-5-phenyl-pentanol*
 2-(2-methylphenyl)-ethanol*
 4-(1-methylethyl)-benzene methanol
 4-(4-hydroxyphenyl)-butan-2-one*
 50 2-phenoxy-ethanol*
 4-(1-methylethyl)-2-hydroxy-1-methyl benzene
 2-methoxy-4-methyl-phenol
 4-methyl-phenol
 anisic alcohol*
 55 p-tolyl alcohol*
 cinnamic alcohol*
 vanillin*
 ethyl vanillin*

eugenol*
 isoeugenol*
 thymol
 anethol*
 5 decahydro 2-naphthalenol
 borneol*
 cedrenol*
 farnesol*
 fenchyl alcohol*
 10 menthol*
 3,7,11-trimethyl-2,6,10-dodecatrien-1-ol
 alpha ionol*
 tetrahydro ionol*
 2-(1,1-dimethylethyl)-cyclohexanol*
 15 3-(1,1-dimethylethyl)-cyclohexanol*
 4-(1,1-dimethylethyl)-cyclohexanol*
 4-isopropyl cyclohexanol
 6,6-dimethyl-bicyclo[3.3.1]hept-2-ene-2-ethanol
 6,6-dimethyl-bicyclo[3.1.1]hept-2-ene-methanol*
 20 p-menth-8-en-3-ol*
 3,3,5-trimethyl-cyclohexanol
 2,4,6-trimethyl-3-cyclohexenyl-methanol*
 4-(1-methylethyl)-cyclohexane-methanol*
 4-(1,1-dimethylethyl)-cyclohexanol
 25 2-(1,1-dimethylethyl)-cyclohexanol
 2,2,6-trimethyl-alpha-propyl cyclohexane propanol*
 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol*
 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)-pent-4-en-2-ol*
 2-ethyl-4(2,2,3-trimethylcyclopentyl-3-enyl)-but-2-en-1-ol*
 30 4-(5,5,6-trimethylbicyclo[2.2.1]hept-2-yl)-cyclohexanol*
 2-(2-methylpropyl)-4-hydroxy-4-methyl-tetrahydropyran*
 2-cyclohexyl propanol*
 2-(1,1-dimethylethyl)-4-methyl-cyclohexanol*
 1-(2-tert-butyl-cyclohexyloxy)-2-butanol*
 35 1-(4-isopropyl-cyclohexyl)-ethanol*
 1-(4-hydroxyphenyl)-butan-3-one
 2,6-dimethyl-oct-7-en-2-ol*
 2,6-dimethyl-heptan-2-ol*
 3,7-dimethyl-octa-1,6-dien-3-ol*
 40

whereby * indicates preferred alcohols and phenols.

[0044] Compounds of the formula I may generate the following organoleptic amines:

anthranilic acid 1-methyl-1-(4-methyl-3-cyclohexen-1-yl)ethyl ester
 45 benzopyrrole
 8,8-di(1H-indol-1-yl)-2,6-dimethyl-octane-2-ol
 anthranilic acid allyl ester
 anthranilic acid 1,5-dimethyl-1-vinyl-4-hexenyl ester
 2-amino-benzoic acid methyl ester*
 50 methyl anthranilic acid N-(2-methylpent-1-en-1-yl) ester
 1-methyl-1-(4-methyl-3-cyclohexen-1-yl)ethyl anthranilic acid
 anthranilic acid phenylethyl ester*
 2-methylamino-benzoic acid methyl ester*
 6-methyltetrahydro-quinoline
 55 isobutyl N-methyl anthranilate
 (Z)-3-hexenyl-2-aminobenzoate*

whereby * indicates the preferred organoleptic amines.

[0045] Compounds of formula I may generate the following organoleptic nitriles:

dodecanenitrile*
 tetradecanenitrile*
 5 5-methyl-7-(1-methylethyl)-bicyclo[2.2.2]oct-5-ene-2-carbonitrile*
 1,2,3,4-tetrahydro-4,4-dimethyl-1-naphthalenecarbonitrile*
 2,4-dimethyl-cyclohex-3-ene-1-nitrile
 2,2,3-trimethyl-3-cyclopentene-1-acetonitrile
 4-(1,1-dimethylethyl)-benzeneacetonitrile
 10 3,12-tridecadienenitrile
 3-methyl-octanenitrile
 3-methyl-dodecanenitrile
 2-methyl-decanenitrile
 decanenitrile
 15 octanenitrile
 4,6,6-trimethyl-heptanenitrile
 isotetradecanenitrile
 3,3-dimethyl-bicyclo[2.2.1]heptane-2-carbonitrile
 octahydro-4,7-methano-indene-5-carbonitrile
 20 3,7,7-trimethyl-4-propylidenenitrile-bicyclo[4.1.0] heptane
 3-hexyloxy-propionitrile
 3-(3,7-dimethyl-oct-2,6-dienyloxy)-propionitrile*
 5-cyano-pentan-2-one*
 3-methyl-5-cyano-pentan-2-one
 25 4-cyano-2,2-dimethyl-butanal
 2-(2-cyanoethyl)-cyclohexanone
 3,7-dimethyl-octa-2,6-diennitrile*
 3-methyl-5-phenyl-pent-2-enenitrile
 3,7-dimethyl-oct-6-enenitrile*
 30 1,2,3,4,5,6,7,8-octahydro-8,8-dimethyl-2-naphthalenecarbonitrile
 3-methyl-2(3)-nonenenitrile
 2-propyl-2-hexenenitrile
 4-methyl-2-propyl-hex-2-enenitrile
 tridec-2-enenitrile
 35 3,7-dimethyl-nona-2,6-diennitrile*
 2,3-dimethyl-2-nonenenitrile
 2-undecanenitrile*
 cyclohexylidene-phenyl-acetonitrile*

40 whereby * indicates the preferred nitriles.

[0046] Compounds of formula I may generate the following preferred organoleptic oximes:

1,5-dimethyl-bicyclo[3.2.1]octan-8-one oxime
 2,4,4,7-tetramethylnona-6,8-dien-3-one oxime
 45 5-methyl-heptan-3-one oxime
 1-bicyclo[2.2.1]hept-5-en-2-yl-ethanone oxime

[0047] It is a matter of course, that it is not possible to give a complete list of the organoleptic and/or antimicrobial aryl and heteroaryl ketones, ketones, aldehydes, alcohols, phenols, amines, nitriles, thiols, oximes and lactones which are generated as a result of the desired cleavage of the ketones of the formula I. The skilled person is, however, quite aware of those substances which provide the desired organoleptic, e.g. fragrance and odour masking and/or antimicrobial effects.

[0048] The compounds of the formula I may preferably be used as sustained release odorants but also as sustained agents to mask or attenuate undesirable odours or to provide additional odours initially present in consumer products, i.e. laundry detergents, fabric softeners, fabric softener sheets, cleaning compositions, swimming pool additives, toiletries and cosmetic products such as shampoo, conditioners and sunscreens. Further applications are sustained antimicrobial agents in the same products.

[0049] The amount required to produce the desired, overall effect varies depending upon the particular compounds

of formula I chosen, the product in which it will be used, and the particular effect desired.

[0050] For example, depending upon the selection and concentration of the compound chosen, when a compound of the formula I is added either singly or as a mixture, e.g. to a laundry product composition at levels ranging from about 0.001 to about 10 % by weight, an odorant, i.e. an odoriferous aryl ketone, ketone, aldehyde, alcohol, phenol, amine, lactone, thiol, oxime, nitrile or two or three of these odoriferous substances in an organoleptically effective amount is/are released when the product is used. These newly formed odorant(s) serve to enhance the odour of the fragrance. Depending on the compound of formula I an antimicrobial agent can be released.

[0051] As is evident from the above compilation of organoleptic substances, a broad range of known odorants can be generated from precursors of the present invention. While manufacturing compositions the precursors of the present invention may be used according to methods known to the perfumer, such as e.g. from W.A. Poucher, *Perfumes, Cosmetics, Soaps*, 2, 7th Edition, Chapman and Hall, London 1974.

[0052] The compounds of formula I can be prepared by using standard methods known to the skilled chemist. A wide variety of methods for their preparation is known.

[0053] Convenient methods are outlined in the Examples without limiting the invention thereto.

Example 1

p-Methoxyphenacyl acetate

[0054] To a solution of 63 g potassium acetate in 750 ml of methanol, 92.9 g of 4-methoxyphenacyl bromide was dropped in. The resulting solution was refluxed for 4 hours, then cooled and poured into water. The aqueous layer was extracted with ether, then the combined organic phases were washed with water, dried and evaporated to dryness. The resulting solid was recrystallized from methanol to yield 54.5 g crystals.

NMR (CDCl₃) δ 8.05-7.70 (m, 2H), 7.10-6.70 (m, 2H), 5.30 (s, H), 3.90 (s, 3H), 2.20 (s, 3H)ppm.

Example 2

p-Tolyloxy-acetic acid 2-oxo-2-phenyl-ethyl ester

[0055] According to the procedure of Example 1, p-tolyloxy-acetic acid 2-oxo-2-phenyl-ethyl ester was prepared from (4-methylphenoxy) acetic acid and 2-bromoacetophenone.

Example 3

2-Amino-3-(3,4-dihydroxy-phenyl)-propionic acid 2-oxo-2-phenyl-ethyl ester

[0056] According to the procedure of Example 1, 2-amino-3-(3,4-dihydroxy-phenyl)-propionic acid 2-oxo-2-phenyl-ethyl ester was prepared from 3-(3,4-dihydroxy phenyl)alanine and 2-bromoacetophenone.

Example 4

Acetic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

[0057] According to the procedure of Example 1, acetic acid 1-methyl-2-oxo-2-phenyl-ethyl ester was prepared from acetic acid and 2-bromo-1-phenyl-1-propan-1-one.

Example 5

Naphthalen-2-yloxy)-acetic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester

[0058] According to the procedure of Example 1, (naphthalen-2-yloxy)-acetic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester was prepared from (2-naphthoxy)acetic acid and 4-methoxy-phenacyl bromide.

Example 6Cyclopropane carboxylic acid 2-naphthalen-2-yl-2-oxo-ethyl ester

- 5 [0059] According to the procedure of Example 1, cyclopropane carboxylic acid 2-naphthalen-2-yl-2-oxo-ethyl ester was prepared from cyclopropane carboxylic acid and 2-bromo-1-naphthalen-2-yl-ethanone.

Example 710 Acetic acid 2-(3-ethyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)-2-oxo-ethyl ester

- [0060] According to the procedure of Example 1, acetic acid 2-(3-ethyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-naphthalen-2-yl)-2-oxo-ethyl ester was prepared from acetic acid and 2-bromo-1-(3-ethyl-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)-ethanone.

15 Example 83-(3-Isopropylphenyl)butanol

- 20 [0061] To a refluxing solution of 132.8 g aluminum isopropoxide in 800 ml of isopropanol, a solution of 300 g 3-(3-isopropylphenyl)butanal in 200 ml of isopropanol was dropped in. The resulting solution was refluxed for 1.5 hours, then cooled down and diluted with ether. Then the organic phase was washed with brine, dried and evaporated to dryness. The resulting oil was distilled to yield 223.5 g of a colourless oil.

- 25 NMR (CDCl₃) δ 7.30-6.96 (m, 4H), 3.69-3.43 (t, 2H), 2.99-2.73 (m, 2H), 1.92-1.77 (m, 2H), 1.60-1.44 (br s, OH), 1.39-1.19 (m, 9H)ppm.

Example 930 Octyloxy-acetonitrile

- [0062] To a suspension of 15.60 g sodium hydride (60%) in 200 ml of 1,2-dimethoxyethane, 50.79 g 1-octanol was dropped in and then the mixture was stirred for 45 min. at room temperature. Then 46.76 g bromoacetonitrile was dropped in, the reaction mixture was refluxed for 2 hours, quenched with water, filtrated and diluted with brine. After
35 extraction with ether, the organic layer was washed with brine and water, dried, filtered and evaporated to dryness. The resulting dark oil was purified by distillation to yield a yellow oil.

NMR (CDCl₃) δ 4.24 (s, 2H), 3.63-3.52 (t, 2H), 1.75-1.50(m, 2H), 1.49-1.15 (m, 10H), 0.96-0.80 (t, 3H)ppm.

40 Example 10Undec-10-enyloxy-acetonitrile

- [0063] According to the procedure of Example 9, undec-10-enyloxy-acetonitrile was prepared from undec-10-en-
45 1-ol, bromoacetonitrile and sodium hydride.

Example 1150 [3-(3-Isopropyl-phenyl)-butoxy]-acetonitrile

- [0064] According to the procedure of Example 9, [3-(3-isopropylphenyl)-butoxy]-acetonitrile was prepared from 3-(3-isopropylphenyl)butanol, bromoacetonitrile and sodium hydride.

55 Example 12[1-Methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allyloxy]-acetonitrile

- [0065] According to the procedure of Example 9, [1-methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allyloxy]-acetonitrile

was prepared from 4-(2,6,6-trimethyl-cyclohex-1-enyl)but-3-en-2-ol (Marvell et al., J. Org. Chem., 37 (1972) 19, 2992-97), bromoacetonitrile and sodium hydride.

Example 13

(2-Pentyl-3-phenyl-allyloxy)-acetonitrile

[0066] According to the procedure of Example 9, (2-pentyl-3-phenyl-allyloxy)-acetonitrile was prepared from amyl cinnamic alcohol, bromoacetonitrile and sodium hydride.

Example 14

2-(3,7-Dimethyl-oct-6-enyloxy)-1-naphthalen-2-yl-ethanone

[0067] To a suspension of 2.24 g magnesium in ether, a solution of 19.09 g 2-bromonaphthalene in 80 ml of ether was dropped in and then the mixture was heated to reflux. After refluxing for one hour, the mixture was cooled down to 0-5°C and a solution of 15.00 g (3,7-dimethyl-oct-6-enyloxy)-acetonitrile (Kulka, K. et al., Perfumer & Flavorist Vol. 3 (1978), p.39) in 20 ml of ether was dropped in at this temperature. After 3 hours stirring at room temperature, the mixture was quenched with water and aqueous sulfuric acid. After extracting with ether, the combined organic phases were washed with saturated sodium bicarbonate and water, dried, filtered and evaporated to dryness. The resulting dark oil was purified by chromatography to yield 15.22 g of a yellow oil.

NMR (CDCl₃) δ 8.49 (s, 1H), 8.05-7.79 (m, 4H), 7.67-7.42 (m, 2H), 5.15-5.01 (t, 1H), 4.83 (s, 2H), 3.73-3.55 (t, 2H), 2.11-1.83, (m, 2H), 1.82-1.03 (m, 11H), 0.96-0.81 (d, 3H)ppm.

Example 15

2-Ethoxy-1-phenyl-ethanone

[0068] According to the procedure of Example 12, 2-ethoxy-1-phenylethanone was prepared from bromobenzene and ethoxyacetonitrile.

Example 16

1-Naphthalen-2-yl-2-octyloxy-ethanone

[0069] According to the procedure of Example 12, 1-naphthalen-2-yl-2-octyloxy-ethanone was prepared from 2-bromonaphthalene and octyloxy-acetonitrile.

Example 17

2-(3,7-Dimethyl-oct-6-enyloxy)-1-(4-methoxy-phenyl)-ethanone

[0070] According to the procedure of Example 12, 2-(3,7-Dimethyl-oct-6-enyloxy)-1-(4-methoxy-phenyl)-ethanone was prepared from 4-bromoanisole and (3,7-dimethyl-oct-6-enyloxy)-acetonitrile (Kulka, K. et al., Perfumer & Flavorist Vol. 3 (1978), p.39).

Example 18

2-(3,7-Dimethyl-oct-6-enyloxy)-1-(4-hydroxy-phenyl)-ethanone

[0071] According to the procedure of Example 12, 2-(3,7-Dimethyl-oct-6-enyloxy)-1-(4-hydroxy-phenyl)-ethanone was prepared from (4-bromo-phenoxy)-trimethyl-silane (J. Org. Chem., 48 (1983), 1543) and (3,7-dimethyl-oct-6-enyloxy)-acetonitrile (Kulka, K. et al., Perfumer & Flavorist Vol. 3 (1978), p.39).

Example 191-Naphthalen-2-yl-2-undec-10-enyloxy-ethanone

5 [0072] According to the procedure of Example 12, 1-naphthalen-2-yl-2-undec-10-enyloxy-ethanone was prepared from 2-bromonaphthalene and undec-10-enyloxy-acetonitrile.

Example 2010 2-[3-(3-Isopropyl-phenyl)-butoxy]-1-naphthalen-2-yl-ethanone

[0073] According to the procedure of Example 12, 2-[3-(3-isopropylphenyl)-butoxy]-1-naphthalen-2-yl-ethanone was prepared from 2-bromonaphthalene and [3-(3-isopropyl-phenyl)-butoxy]acetonitrile.

15 Example 212-[3-(3-Isopropyl-phenyl)-butoxy]-1-(4-methoxy-phenyl)-ethanone

20 [0074] According to the procedure of Example 12, 2-[3-(3-isopropyl-phenyl)-butoxy]-1-(4-methoxy-phenyl)-ethanone was prepared from 4-bromoanisole and [3-(3-isopropyl-phenyl)-butoxy]-acetonitrile.

Example 2225 2-(3,7-Dimethyl-oct-6-enyloxy)-1-phenantren-9-yl-ethanone

[0075] According to the procedure of Example 12, 2-(3,7-dimethyl-oct-6-enyloxy)-1-phenantren-9-yl-ethanone was prepared from 9-bromo-phenantrene and (3,7-dimethyl-oct-6-enyloxy)-acetonitrile (Kulka, K. et al., Perfumer & Flavorist Vol. 3 (1978), p.39).

30 Example 231-(4-Methoxy-phenyl)-2-[1-methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allyloxy]-ethanone

35 [0076] According to the procedure of Example 12, 1-(4-methoxy-phenyl)-2-[1-methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allyloxy]-ethanone was prepared from 4-bromoanisole and [1-methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allyloxy]-acetonitrile.

Example 2440 1-(4-Methoxy-phenyl)-2-(2-pentyl-3-phenyl-allyloxy)-ethanone

[0077] According to the procedure of Example 12, 1-(4-methoxy-phenyl)-2-(2-pentyl-3-phenyl-allyloxy)-ethanone was prepared from 4-bromoanisole and (2-pentyl-3-phenyl-allyloxy)-acetonitrile.

45 Example 25Hydroxy-acetic acid 2-oxo-2-phenyl-ethyl ester

50 [0078] A mixture of 7.61 g glycolic acid, 19.90 g phenacylbromide and 13.9 ml triethylamine in 300 ml of ethyl acetate was stirred for 20 hours at room temperature. Then the mixture was filtered and the filtrate was washed with 2N HCl, saturated sodium bicarbonate and brine, dried, filtered and evaporated to dryness. The resulting white solid was purified by recrystallisation to yield 11.85 g of colourless crystals.

NMR (CDCl₃) δ 7.97-7.86 (m, 2H), 7.71-7.44 (m, 3H), 5.49 (s, 2H), 4.45-4.36 (d, 2H), 2.59-2.47 (t, 1H)ppm.

55

Example 26Carbonic acid 4-allyl-2-methoxy-phenyl ester 2-(4-methoxy-phenyl)-2-oxo-ethyl ester

[0079] To a solution of 17.62 g 2-hydroxy-4'-methoxyacetophenone (Kovach et al., J. Amer. Chem. Soc., Vol. 115, No. 23, (1993) p. 10476) and 12.96 g pyridine in 240 ml of dichloromethane, a solution of 26.32 g eugenol-chloroformate in 60 ml of dichloromethane was dropped in at 5-10°C and then the mixture was stirred for 6 hours at room temperature. Then the reaction mixture was acidified with 2N HCl and extracted with ether. The combined organic phases were washed with 2N HCl, saturated sodium bicarbonate and brine, dried, filtered and evaporated to dryness. The resulting solid was purified by recrystallisation to yield 29.79 g of yellow crystals.

NMR (CDCl₃) δ 7.98-7.86 (m, 2H), 7.20-6.72 (m, 5H), 6.09-5.83 (m, 1H), 5.41 (s, 2H), 5.18-5.02 (m, 2H), 3.89 (s, 3H), 3.87 (s, 3H), 3.43-3.32 (d, 2H) ppm.

Example 27Carbonic acid 3-methyl-5-phenyl-pentyl ester 2-naphthalen-2-yl-2-oxo-ethyl ester

[0080] According to the procedure of Example 22, carbonic acid 3-methyl-5-phenyl-pentyl ester 2-naphthalen-2-yl-2-oxo-ethyl ester was prepared from 2-hydroxy-1-naphthalen-2-yl-ethanone (Langenbeck et al., Chem. Ber., 69 (1936) 514-516), 3-methyl-5-phenyl-pentanol-chloroformate and pyridine.

Example 284-Hydroxy-undecanoic acid sodium salt

[0081] To a solution of 43.6 g sodium hydroxide in 150 ml of methanol heated to reflux, 200 g gamma-undecalactone were dropped in. After stirring 2 hours at reflux, the mixture was cooled to room temperature and evaporated to dryness. The resulting crystals were washed with hexane to yield 240 g of colourless crystals.

NMR (CDCl₃) δ 5.1-4.8 (br s, OH), 3.63-3.42 (m, 1H), 2.39-2.20 (t, 2H), 1.89-1.52 (m, 2H), 1.51-1.15 (m, 12H), 1.00-0.81 (t, 3H) ppm.

Example 294-Hydroxy-undecanoic acid-2-naphthalen-2-yl-2-oxo-ethyl ester

[0082] A suspension of 20.00 g 2-bromo-2'-acetonaphthone, 18.01 g 4-hydroxy-undecanoic acid sodium salt and 0.5 g tetrabutylammonium bromide in 180 ml of dimethylformamide was stirred for 6 hours at 50°C. Then the reaction mixture was diluted with ether and washed with water, 2N HCl, saturated sodium bicarbonate and brine, dried, filtered and evaporated to dryness. The resulting solid was purified by recrystallisation to yield 7.92 g of colourless crystals.

NMR (CDCl₃) δ 8.41 (s, 1H), 8.03-7.78 (m, 4H), 7.70-7.48 (m, 2H), 5.55-5.42 (2s, 2H), 3.80-3.61 (m, 1H), 2.27-2.60 (t, 2H), 2.33-2.06 (br s, OH), 2.05-1.11 (m, 14H), 1.02-0.79 (t, 3H) ppm.

Example 304-Hydroxy-undecanoic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester

[0083] According to the procedure of Example 24, 4-hydroxy-undecanoic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester was prepared from 2-bromo-4'-methoxyacetophenone, 4-hydroxy-undecanoic acid sodium salt and tetrabutylammonium bromide.

Example 314-Hydroxy-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

[0084] According to the procedure of Example 24, 4-hydroxy-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

was prepared from 2-bromopropiophenone, 4-hydroxy-undecanoic acid sodium salt and tetrabutylammonium bromide.

Example 32

4-(3-Methyl-5-phenyl-pentyloxycarbonyloxy)-undecanoic acid 2-naphthalen-2-yl-2-oxo-ethyl ester

[0085] To a solution of 4.00 g 4-hydroxy-undecanoic acid-2-naphthalen-2-yl-2-oxo-ethyl ester and 1.71 g pyridine in 20 ml of THF, a solution of 2.91 g 3-methyl-5-phenyl-pentanol-chloroformate in 10 ml of THF was dropped in at 0°C and then the mixture was stirred for 3 hours at room temperature. Then the reaction mixture was quenched with water and extracted with ether. The organic phase was washed with 2N HCl, saturated sodium bicarbonate and water, dried, filtered and evaporated to dryness. The resulting yellow oil was purified by chromatography to yield 5.46 g of a colorless oil.

NMR (CDCl₃) δ 8.42 (s, 1H), 8.03-7.80 (m, 4H), 7.70-7.45 (m, 2H), 7.35-7.03 (m, 5H), 5.49 (s, 2H), 4.89-4.71 (m, 1H), 4.30-4.07 (m, 2H), 2.80-2.45 (m, 4H), 2.21-1.12 (m, 19H), 1.10-0.78 (m, 6H) ppm.

Example 33

4-Hex-3-enyloxycarbonyloxy-undecanoic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester

[0086] According to the procedure of Example 26, 4-hex-3-enyloxycarbonyloxy-undecanoic acid 2-(4-methoxy-phenyl)-2-oxoethyl ester was prepared from 4-hydroxy-undecanoic acid 2-(4-methoxy-phenyl)-2-oxo-ethyl ester, cis-hex-3-enol-chloroformate and pyridine.

Example 34

4-(3-Methyl-5-phenyl-pentyloxycarbonyloxy)-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

[0087] According to the procedure of Example 26, 4-(3-methyl-5-phenyl-pentyloxycarbonyloxy)-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester was prepared from 4-hydroxy-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester, 3-methyl-5-phenyl-pentanol-chloroformate and pyridine.

Example 35

4-[2-Ethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-2-enyloxycarbonyloxy]-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

[0088] According to the procedure of Example 26, 4-[2-ethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-2-enyloxycarbonyloxy]-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester was prepared from 4-hydroxy-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester, 2-ethyl-4-(2,2,3-trimethyl-cyclopentyl-3-en-1-yl)-but-2-en-1-ol chloroformate and pyridine.

Example 36

4-[1,2-Dimethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-3-enyloxycarbonyloxy]-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester

[0089] According to the procedure of Example 26, 4-[1,2-dimethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-3-enyloxycarbonyloxy]-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester was prepared from 4-hydroxy-undecanoic acid 1-methyl-2-oxo-2-phenyl-ethyl ester, 3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-ol chloroformate and pyridine.

Example 37

[0090] Test cloth was washed with detergent to which one or more of the precursors of Examples 1-7, 12-22, 24-26 had been added. The cloth was then line dried. The cloth dried in sunlight had a distinct fragrance note, as determined by a trained panel. In contrast, the cloth dried without light was olfactively neutral.

Example 38

[0091] Test cloth was washed with a detergent and then a fabric softener, containing one or more of the precursors of Examples 1-7, 12-22, 24-26 was added to the rinse cycle. The cloth was then line dried. The cloth dried in sunlight had a distinct fragrance note, as determined by a trained panel. In contrast, the cloth dried without sunlight was olfactively neutral.

Example 39Preparation of an O/W sunscreen lotion UV-B and UV-A:

[0092] Sunscreen lotion containing 0.5 % of one or more compounds of Examples 1-7, 12-22, 24-26.

Recipe: %	compound
Part A:	
2 %	Octyl methoxycinnamate
3 %	4-4-Butyl-4'methoxydibenzoyl methane
12 %	Coco-caprylate/caprate
4 %	Isostearyl neopentanoate
0.25 %	Diethyleneglycol monostearate
1 %	Cetylalcohol
0.25 %	Methyl-propylparabene
0.1 %	EDTA-sodium salt
1 %	Diethanolamine cetylphosphate
Part B:	
20 %	Acrylate C10-C30 Alkylacrylate
50.1 %	water deion.
5 %	1,2-Propanediol
0.8 %	Potassium hydroxide.
Part A was heated in a reactor to 85°C.	
Part B was slowly added within 10 min., followed by addition of KOH and 0.5 % of one or more of the compounds in Examples 1-7, 12-22, 24-26. The emulsion was then cooled and degassed.	

Example 40

[0093]

a) Fabric softener of the ester quat type (4 x concentrate):

INGREDIENTS	CHEMICAL NAME	%
PHASE A		
DEIONISED WATER		to 100.0
MgCl ₂ (saturated sol.)	Magnesium chloride	1.0
PHASE B		
REWOQUAT WE 18Di-	(tallow carboxyethyl)hydroxy ethyl methylammonium methosulfate	15.0
GENAPOL O 100	Ethoxylated fatty alcohol C16-C18 10EO	2.0
ANTIFOAM DB 31		0.5

(continued)

INGREDIENTS	CHEMICAL NAME	%
<u>PHASE B</u>		
<u>PHASE C</u>		
ISOPROPYL ALCOHOL		3.0
PRESERVATIVE		Qs
PERFUME		Qs

PROCESS:

While stirring and heating to 65° C, mix part A, then part B preheated to 65° C. After cooling to room temperature, add part C. The pH value of the finished product is 2.60. Recommended level of perfume is 1.0 %. Delayed release fragrances of Examples 1-7, 12-22, 24-26 may be any part of this 1.0 %.

b) Fabric softener of the ester quat type (1 x concentrate):

INGREDIENTS	CHEMICAL NAME	%
<u>PHASE A</u>		
DEIONISED WATER		to 100.0
<u>PHASE B</u>		
REWOQUAT WE 18Di-(tallow carboxyethyl)hydroxy ethyl methylammonium methosulfate		6.0
DOBANOL 25-9Ethoxylated fatty alcohol C12-C15 9EO		0.50
ANTIFOAM DB 31		0.10
<u>PHASE C</u>		
MYACIDE BT 302-bromo-2-nitropropane 1,3 diol		0.03
PROXEL GXLBenzisothiazolinone sodium salt		0.02
PERFUME		Qs


PROCESS:

While stirring and heating to 65° C, mix part A, then part B preheated to 65° C. After cooling to room temperature, add part C. The pH value of the finished product is 3.50.

Recommended level of perfume: 0.3 %. Delayed release fragrances of Examples 1-7, 12-22, 24-26 may be any part of this 0.3 %.

Claims

1. Ketones of the formula I



wherein

Y is an optionally substituted alkyl, cycloalkyl, cycloalkylalkyl, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, alkoxy carbonyl, cycloalkoxy carbonyl, aryloxy carbonyl, amide, oximinocarbonyl, alkylthiocarbonyl, cycloalkylthiocarbonyl, whereby all alkyl groups may be straight or branched and all alkyl and cycloalkyl groups may be saturated or unsaturated,

R¹ is hydrogen or a C₁₋₆ alkyl group which may be substituted, saturated or unsaturated, straight or branched,

A is a chromophoric aromatic or heteroaromatic ring or ring system,

A may carry 1 to 8 substituents selected from hydrogen, hydroxy, mercapto, amino, acyl, ester, esterified hydroxy group, cyano and nitro; alkyl, alkoxy, alkylthio and arylalkoxy which may be substituted, saturated or unsaturated, straight or branched; aryl and heteroaryl which may be substituted; one of the substituents may be a polymeric group, and

n is an integer


with the exception of

2-ethoxy-1-phenyl-ethanone,

hydroxy-acetic acid 2-oxo-2-phenyl-ethyl ester,


acetic acid 1-methyl-2-oxo-2-phenyl-ethyl ester, and p-methoxyphenacyl acetate.

2. Ketones according to claim 1, which yield after cleavage by light one or more organoleptic compounds.
3. Ketones according to claim 1 or 2, which yield after cleavage by heat one or more organoleptic compounds.
4. Ketones according to any of claims 1 to 3, which yield after cleavage by hydrolysis one or more organoleptic compounds.
5. Ketones according to any of claims 1 to 4, which yield after cleavage by enzymes one or more organoleptic compounds.
6. Ketones according to any of claims 1 to 5, which yield after cleavage two or more organoleptic compounds.
7. Ketones according to any of claims 1 to 6, which yield after cleavage an organoleptic aryl ketone of the formula II




8. Ketones according to any of claims 1 to 7, which yield after cleavage one or more organoleptic aldehydes or ketones derived from O-Y.
9. Ketones according to any of claims 1 to 7, which yield an organoleptic alcohol, phenol, aldehyde or keton derived from Y.

10. Ketones according to any of claims 1 to 7, which yield after cleavage an organoleptic lactone derived from Y.
11. Ketones according to any of claims 1 to 7, which yield after cleavage an amine used in perfumery and derived from Y.
12. Ketones according to any of claims 1 to 11, wherein R¹ is hydrogen or methyl.
13. Ketones according to any of claims 1 to 12, wherein R¹ is hydrogen or methyl and A carries no more than three substituents, preferably methyl groups.
14. Ketones according to any of claims 1 to 12, wherein R¹ is hydrogen and A is one aromatic ring which is substituted by one methoxy group, preferably in the para-position.
15. Ketones according to any of claims 1 to 13, wherein A is a ring system of the formula



and both rings may be substituted preferably with one or more methyl groups.

16. Ketones according to any of claims 1 to 15, wherein n is 1 or 2.
17. Ketones according to any of claims 1 to 16, wherein one of the substituents of A is a polymeric group.
18. Ketones to any of the claims 1 to 17, wherein the groups for which Y stands are substituted with one or more substituents selected from aryl; cycloalkyl; cycloalkenyl; oxo; oxy; cyano; amino; hydroxy; hydroxy being esterified with a carboxylic acid, a carbonic acid ester (carbonate), a N,N-substituted carbamic acid (carbamate) or an oximino carbonic acid; amino being substituted with oximinocarbonyl; an esterified carboxylic group; and a carboxylic group esterified with an oxime.
19. Use of ketones of formula I



wherein

Y is an optionally substituted alkyl, cycloalkyl, cycloalkylalkyl, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, alkyloxycarbonyl, cycloalkyloxycarbonyl, aryloxycarbonyl, amide, oximinocarbonyl, alkylthiocarbonyl, cycloalkylthiocarbonyl, whereby all alkyl groups may be straight or branched and all alkyl and cycloalkyl groups may be saturated or unsaturated,

R¹ is hydrogen or a C₁₋₆ alkyl group which may be substituted, saturated or unsaturated, straight or branched,

A is a chromophoric aromatic or heteroaromatic ring or ring system,

A may carry 1 to 8 substituents selected from hydrogen, hydroxy, mercapto, amino, acyl, ester, esterified hydroxy group, cyano and nitro; alkyl, alkoxy, alkylthio and arylalkoxy which may be substituted, saturated or

unsaturated, straight or branched; aryl and heteroaryl which may be substituted; one of the substituents may be a polymeric group, and

n is an integer,

as precursors.

20. Use according to claim 19 as precursors for fragrances and/or antimicrobial agents.

21. Use according to claim 20 in detergents, fabric softeners, personal care products, cleaning compositions, cosmetics and sun protection products.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 983 990 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
02.08.2000 Bulletin 2000/31

(43) Date of publication A2:
08.03.2000 Bulletin 2000/10

(21) Application number: **99810685.0**

(22) Date of filing: **29.07.1999**

(51) Int Cl.7: **C07C 49/84, C07C 69/145,
C07C 69/712, C07C 69/74,
C07C 69/675, C07C 69/24,
C07C 69/96, C07C 229/22,
C07C 329/06, C11D 3/50,
C11D 3/20, A61K 7/46**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **04.09.1998 EP 98810883**

(71) Applicant: **Glvaudan Roure (International) S.A.**
1214 Vernier-Genève (CH)

(72) Inventors:
• **Anderson, Denise Dr.**
8032 Zürich (CH)
• **Fráter, Georg Dr.**
8400 Winterthur (CH)

(74) Representative: **Patentanwälte**
Schaad, Balass, Menzi & Partner AG
Dufourstrasse 101
Postfach
8034 Zürich (CH)

(54) **Ketones useful as precursors for organoleptic compounds**

(57) The ketones of the formula I are useful for the delivery of organoleptic compounds, especially for fla-

vours, fragrances and masking agents and antimicrobial compounds.

EP 0 983 990 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 99 81 0685

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	CHEMICAL ABSTRACTS, vol. 108, no. 3, 18 January 1988 (1988-01-18) Columbus, Ohio, US; abstract no. 021462, MAHAJAN S D ET AL: "Polymer-supported reactions. Synthesis of phenacyl esters of phenoxyacetic acids" XP002139246 * abstract * & INDIAN J. CHEM., SECT. B (IJSBDB,03764699);1987; VOL.26B (2); PP.186, Shivaji Univ.;Dep. Chem.; Kolhapur; 416 004; India (IN)	1	C07C49/84 C07C69/145 C07C69/712 C07C69/74 C07C69/675 C07C69/24 C07C69/96 C07C229/22 C07C329/06 C11D3/50 C11D3/20 A61K7/46
A	WO 98 07405 A (PROCTER & GAMBLE) 26 February 1998 (1998-02-26) * claims *	1-21	
D,A	US 5 726 345 A (PAGET WALTER ET AL) 10 March 1998 (1998-03-10) * the whole document *	1-21	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07C C11D
INCOMPLETE SEARCH <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		31 May 2000	BONNEVILLE, E
CATEGORY OF CITED DOCUMENTS <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 (03.82) (P04C27)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 99 81 0685

Claim(s) searched incompletely:
1-21

Reason for the limitation of the search:

The search revealed such a large number of particularly relevant documents, in particular with regard to novelty, that the drafting of a comprehensive European Search Report is not feasible. The cited documents are considered as to form a representative sample of the revealed documents, duly taking into account their relevance with respect to the subject-matter as illustrated by the examples.

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 99 81 0685

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D, A	EP 0 816 322 A (GIVAUDAN ROURE INT) 7 January 1998 (1998-01-07) * the whole document *	1-21	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 81 0685

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

31-05-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9807405 A	26-02-1998	AU 3916697 A	06-03-1998
		AU 3985397 A	06-03-1998
		AU 3985597 A	06-03-1998
		AU 3985997 A	06-03-1998
		BR 9712787 A	14-12-1999
		CN 1233281 A	27-10-1999
		CN 1233282 A	27-10-1999
		CN 1233283 A	27-10-1999
		CN 1233284 A	27-10-1999
		CN 1233947 A	03-11-1999
		CZ 9900541 A	14-07-1999
		CZ 9900563 A	14-07-1999
		CZ 9900564 A	11-08-1999
		EP 0951274 A	27-10-1999
		EP 0955994 A	17-11-1999
		EP 0927238 A	07-07-1999
		EP 0921824 A	16-06-1999
		EP 0922083 A	16-06-1999
		EP 0922084 A	16-06-1999
		EP 0922085 A	16-06-1999
		JP 2000501450 T	08-02-2000
		JP 2000502746 T	07-03-2000
		JP 2000502400 T	29-02-2000
		WO 9807407 A	26-02-1998
		WO 9807809 A	26-02-1998
		WO 9807810 A	26-02-1998
		WO 9807811 A	26-02-1998
		WO 9807683 A	26-02-1998
		WO 9807455 A	26-02-1998
		WO 9807812 A	26-02-1998
		WO 9807813 A	26-02-1998
		WO 9807814 A	26-02-1998
US 5726345 A	10-03-1998	AU 684804 B	08-01-1998
		AU 7194094 A	28-02-1995
		CN 1113388 A,B	13-12-1995
		EP 0668904 A	30-08-1995
		WO 9504809 A	16-02-1995
		JP 8502522 T	19-03-1996
		US 5649979 A	22-07-1997
EP 0816322 A	07-01-1998	BR 9405545 A	25-05-1999
		EP 0815816 A	07-01-1998
		AU 3221697 A	21-01-1998
		CA 2259398 A	08-01-1998
		JP 11513919 T	30-11-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

